



Olgoonik
Fairweather LLC



Temporal variability of biodiversity in Arctic and subArctic marine benthic communities

Arny L. Blanchard
Institute of Marine Science
University of Alaska Fairbanks

Alaska Marine Science Symposium, January 20-22, 2014, Anchorage, Alaska



Acknowledgments

- The Port Valdez Environmental Studies Program was funded by Alyeska Pipeline Service Co.
 - I thank the many technicians and staff who assisted, Howard Feder who initiated the study, the crew of the Valdez Star, and Dave Connor of Alyeska for his support.
- The Chukchi Sea Environmental Studies Program was funded by
 - I thank ConocoPhillips Company, Shell Exploration & Production Company, and Statoil USA E&P, Inc. for funding the study, the many technicians and staff who assisted and crews of the Bluefin and Westward Wind who assisted us, and Aldrich Offshore Services and Olgoonik Oilfield Services provided logistics and support. Thanks to Sheyna Wisdom for herding the “clams”.


Questions about biodiversity

- Do we know enough about biodiversity?
 - We study distributions of dominants!
- Are spatially disperse stations across heterogeneous habitats really describing biodiversity patterns?
- Do we know enough to draw inferences about marine invaders into the Arctic?

The Port Valdez Environmental Studies Program

PVESP: the first integrated multi-disciplinary environmental study in Alaska, 1971 to present.


- Port Valdez is a glacial fjord, Prince William Sound, Alaska.
- Glacial sediments drive spatial gradients in biota.



The Chukchi Sea Environmental Studies Program


CESP: An integrated multidisciplinary investigation of the northeastern Chukchi Sea ecosystem, 2008-2013.

- Three study areas were sampled in 2008-2013.
- A larger area encompassing Hanna Shoal was sampled 2011-2012.



Spatial patterns of biodiversity



- Environmental characteristics drive spatial patterns in biodiversity.





Spatial patterns of biodiversity

These animals less abundant in finer sediments.

Sternaspidae





Trichobranchidae





Muddy
sediments.

These animals are abundant in fine
sediments.

Paraonidae



Thyasiridae



Silty
sediments.

Sedimentation rate

- Environmental covariates often confused with drivers!

Problems with biodiversity studies


- Species are captured with varying probabilities specific to habitats.
- Species are so poorly known in Alaska that inferences are difficult to make about changes.
- Scales of natural change unknown!

Temporal drivers of diversity


- Water circulation patterns covarying with climate variations can influence benthic fauna observed diversity.

- Port Valdez, Alaska.
- San Francisco Bay, CA.

Pacific Decadal Oscillation





Source: jisao.washington.edu.




Temporal changes in subArctic biodiversity

- Density and richness in Port Valdez moderately to strongly correlated with the PDO.




The Arctic Oscillation

- Higher sea level pressure in the Arctic Ocean (neg. AO phase) results in stronger Beaufort Gyre.





Source: NSIDC.



Source: Morison et al., 2012. Nature.


Temporal changes in Arctic biodiversity

- Strong association between Chukchi Sea subtidal benthic macrofaunal abundance and the Arctic Oscillation (AO; an index of Arctic Ocean sea-level pressure).



Measured biodiversity

>10 years
sampling
needed to
document
richness at
small (PVESP)
and mid-scale
(CSESP)
extents.



Take home message

- Repeated sampling of sites within homogenous systems requires >10 years to begin to understand richness or measured biodiversity.
- Conclusions of biodiversity confounded in short term studies due to temporal variability.
- Temporal drivers must be considered!
 - Might sample at a “peak” or “valley” of an alternate state.